

Phase Separation in the Hubbard Model

Chi Au and Bao-Heng Zhao*

Department of Applied Mathematics

The Hong Kong Polytechnic University, Hong Kong

H. T. Nieh[†]

The Hong Kong University of Science and Technology, Hong Kong

Abstract

Making use of symmetry properties and known exact results for the Hubbard Model on bipartite lattice, we show that (1) there is no phase separation for repulsive coupling at low dopings, (2) phase separation and superconductivity co-exist in the ground states for attractive coupling in a range of filling fractions.

PACS numbers: 71.27.+a, 74.80.-g, 74.25.Bt

*Permanent address: Department of Physics, Graduate School, Chinese Academy of Sciences, P. O. Box 3908, Beijing 100039, China

[†]On leave of absence from Institute for Theoretical Physics, State University of New York at Stony Brook, Stony Brook, NY 11794, USA

Recent experiments indicate the presence of phase separation in cuprate superconductors, namely, there exist separate hole-rich and hole-poor phases[1]-[4]. There have been theoretical studies, all employing approximations or numerical estimations, on this phenomenon based on the t - J model[5]-[6], as well as directly on the Hubbard model[7]-[8]. Most of these studies indicate that these models can explain the phenomenon of phase separation. The exceptions are those of Putikka *et al.*[6], and Moreo *et al.* [7]. While the quantum Monte Carlo calculations of small 2D lattice systems indicate no phase separation for the 2D Hubbard model[7], the investigation using high-temperature expansions for the 2D t - J model suggests the existence of phase separation only for a limited range of J/t values[6].

In this note, we look again at this phenomenon on the basis of the Hubbard model with repulsive on-site interaction. We make use of some exact results for the Hubbard model on bipartite lattice with attractive coupling[9]-[10], which, after applying the well-known unitary particle-hole transformation[11], imply a certain general property for the repulsive coupling case. It in turn implies that the repulsive Hubbard model on bipartite lattice can not accommodate phase separation at low dopings. We also look at the case of attractive Hubbard model, for which we show that, at temperature $T = 0$, phase separation is possible, and actually co-exists with superconductivity.

The Hamiltonian of the Hubbard model on bipartite lattice is

$$H = - \sum_{ij,\sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} + U \sum_i c_{i\uparrow}^\dagger c_{i\uparrow} c_{i\downarrow}^\dagger c_{i\downarrow} - \mu \sum_i (c_{i\uparrow}^\dagger c_{i\uparrow} + c_{i\downarrow}^\dagger c_{i\downarrow}), \quad (1)$$

with the usual notations. The bipartite lattice sites Λ are divided into subsets

A and B . The number of sites of A (B) is denoted by $|A|$ ($|B|$), and the total number of sites by $|\Lambda|$. In (1), t_{ij} vanishes if both i and j belong to the same subset. An exact result obtained by Lieb[9] states that, for attractive coupling ($U < 0$) and even number of electrons, the ground state of the system is unique and has total spin $S = 0$. Thus, at temperature $T = 0$, the system is non-ferromagnetic. This result was extended to $T > 0$ by Kubo and Kishi[10], who find that there is an upper bound on a certain two-point function. It implies absence of long-range magnetic order in attractive Hubbard model at finite temperature.

We introduce an external magnetic field in the z -direction. The Hamiltonian (1) becomes

$$H = - \sum_{ij,\sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} + U \sum_i c_{i\uparrow}^\dagger c_{i\uparrow} c_{i\downarrow}^\dagger c_{i\downarrow} - \mu \sum_i (c_{i\uparrow}^\dagger c_{i\uparrow} + c_{i\downarrow}^\dagger c_{i\downarrow}) + h \sum_i (c_{i\uparrow}^\dagger c_{i\uparrow} - c_{i\downarrow}^\dagger c_{i\downarrow}). \quad (2)$$

Define

$$s^z = \frac{\sum_i s_i^z}{|\Lambda|}, \quad (3)$$

where s_i^z is the z -component of the spin of the particle on site i . In general, we have

$$\lim_{|\Lambda| \rightarrow \infty} \langle s^z \rangle = f(h, T), \quad (4)$$

where $\langle \dots \rangle$ is the average over a grand canonical ensemble. Invoking Lieb-Kubo-Kishi's result, we have

$$f(0, T) = 0, \quad (5)$$

and, furthermore, since there is no long-range magnetic order, that $f(h, T)$ is analytic at $h = 0$. As a consequence, we can draw the conclusion that

$f(h, T)$ is a well-behaved, i. e. continuous and single-valued, function in the neighborhood of $h = 0$ for $T \geq 0$. For small h ,

$$\lim_{|\Lambda| \rightarrow \infty} \langle s^z \rangle = \chi(T)h, \quad (6)$$

where $\chi(T)$ is the spin susceptibility.

We now make the unitary particle-hole transformation[11]

$$c_{i\uparrow} \rightarrow c_{i\uparrow}, \quad c_{i\downarrow} \rightarrow \epsilon_i c_{i\downarrow}^\dagger, \quad (7)$$

where $\epsilon_i = 1$ (-1) on the lattice set A (B). The Hamiltonian (2), up to a constant, is transformed into

$$\begin{aligned} \tilde{H} = & - \sum_{\langle ij \rangle, \sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} + U' \sum_i c_{i\uparrow}^\dagger c_{i\uparrow} c_{i\downarrow}^\dagger c_{i\downarrow} - \mu' \sum_i (c_{i\uparrow}^\dagger c_{i\uparrow} + c_{i\downarrow}^\dagger c_{i\downarrow}) \\ & + h' \sum_i (c_{i\uparrow}^\dagger c_{i\uparrow} - c_{i\downarrow}^\dagger c_{i\downarrow}), \end{aligned} \quad (8)$$

with

$$\begin{aligned} U' &= -U, \\ \frac{U'}{2} - \mu' &= h, \\ h' &= \frac{U}{2} - \mu, \end{aligned} \quad (9)$$

and (6) into

$$\lim_{|\Lambda| \rightarrow \infty} D = 1 + \chi(T)(U' - 2\mu'), \quad (10)$$

where D is the density for the system described by $\tilde{H}$,

$$D = \frac{\sum_{i,\sigma} \langle c_{i\sigma}^\dagger c_{i\sigma} \rangle}{|\Lambda|}. \quad (11)$$

Recall that $U' = -U > 0$.

From (10), we see that for a system described by a repulsive Hubbard model ($U' > 0$) on bipartite lattice, the density D in the thermodynamic

limit is a well-behaved, i. e. *continuous and single-valued*, function of the chemical potential μ' near $\frac{U'}{2}$, namely at low dopings. This behavior is in contrast with what is expected of a first-order phase transition.

When a system with chemical potential μ_o undergoes a first-order phase transition at temperature T , and separates into two phases with different densities, we have

$$\begin{aligned} D_1 &= \lim_{\mu \rightarrow \mu_o + 0} \lim_{|\Lambda| \rightarrow \infty} D, \\ D_2 &= \lim_{\mu \rightarrow \mu_o - 0} \lim_{|\Lambda| \rightarrow \infty} D, \end{aligned} \tag{12}$$

with $D_1 \neq D_2$. The density D , as a function of the chemical potential, is discontinuous at μ_o in the thermodynamic limit when there is first-order phase transition. Thus the continuity of $\lim_{|\Lambda| \rightarrow \infty} D$ in (10) implies that *a repulsive Hubbard model on bipartite lattice can not accommodate co-existence of two phases at low dopings*. This result is valid for general h' , including $h' = 0$, which is the case of vanishing external magnetic field.

The following are two remarks:

1. The conclusion we made with regard to the phase separation in a repulsive Hubbard model on bipartite lattice applies to low dopings. This is because the exact result of Lieb-Kubo-Kishi is for the case of vanishing magnetic field ($h = 0$). However, if the function $f(h, T)$ is also well behaved, as may be reasonably expected for a paramagnetic system, at large values of h , then our conclusion about phase separation can be extended to high dopings.

2. Lieb has proved[9] in the case of repulsive Hubbard model on bipartite lattice with half-filling and $h = 0$ that the (degenerate) ground states have

the same spin $S = \frac{1}{2}(|A| - |B|)$ (if $|A| \geq |B|$), with

$$S_z = -\frac{1}{2}(|A| - |B|), -\frac{1}{2}(|A| - |B|) + 1, \dots, \frac{1}{2}(|A| - |B|) \quad (13)$$

This result can be used to make certain statement concerning phase separation at $T = 0$ for attractive Hubbard model on bipartite lattice. Making use again of the unitary particle-hole transformation, it is straightforward to obtain, for the attractive case, that at $T = 0$,

$$D = 1 - \frac{|A| - |B|}{2|\Lambda|}, 1 - (\frac{|A| - |B|}{2|\Lambda|} - \frac{1}{|\Lambda|}), \dots, 1 + \frac{|A| - |B|}{2|\Lambda|}. \quad (14)$$

This implies that in an attractive Hubbard model on bipartite lattice different phases can co-exist at $T = 0$, with the densities having values given in (14).

It has been shown by Shen and Qiu[12] that for attractive Hubbard model on bipartite lattice with $|A| - |B| = O(|\Lambda|)$, the ground states with densities in the range $(1 - \frac{|A| - |B|}{2|\Lambda|}) < D < (1 + \frac{|A| - |B|}{2|\Lambda|})$ have off-diagonal long-range order (ODLRO) in the two-particle reduced density matrix, and, consequently, posses the property of superconductivity[13]. That is, superconductivity and phase separation co-exist in this case.

In conclusion, we have investigated the question of phase separation for the Hubbard model on bipartite lattice. Our result that there is no phase separation at low dopings for $T \geq 0$, being based on the exact results of Lieb[9], and Kubo-Kishi[10], is an *exact* statement of the repulsive Hubbard model on bipartite lattice of any dimension. It is consistent with the quantum Monte Carlo calculations of small 2D lattice systems for the Hubbard Model[7]. It is also consistent with the high-temperature expansion result of Putikka *et al.*[6] for the 2D t - J model, to the extent that the Hubbard model and the t - J model are expected to have similar results at small J/t values.

Acknowledgment

We acknowledge discussions with Prof. Fu-Chun Zhang, and thank him for pointing out the work of Putikka et al.[6] This work is supported in part by The Hong Kong Polytechnic University. One of the authors (B. H. Zhao) is grateful to the hospitality of the Department of Applied Mathematics, The Hong Kong Polytechnic University. He is also supported in part by NNSF of China.

References

- [1] P. C. Hammel, A. P. Reyes, Z. Fisk, M. Takigawa, J. D. Thompson, R. H. Heffner, S-W. Cheong, and J. E. Shirber, Phys. Rev. B **42**, 6781 (1990).
- [2] R. K. Kremer, E. Sigmund, V. Hizhnyakov, F. Hentsch, A. Simon, K. A. Müller, and M. Mehring, Z. Phys. B - Condensed Matter **86**, 319 (1992).
- [3] *Proceedings of the Workshop on Phase Separation in Cuprate Superconductors*, edited by K. A. Müller and G. Benedek (World Scientific, Singapore, 1993).
- [4] *Phase Separation in Cuprate Superconductors*, edited by E. Sigmund and K. A. Müller (Springer-Verlag, Berlin Heidelberg, 1994).
- [5] M. Marder, N. Papanicolaou, and G. C. Psaltakis, Phys. Rev. B **41**, 6920 (1990).

- S. A. Kivelson, V. J. Emery, and H. Q. Lin, Phys. Rev. B **42**, 6523 (1990).
- A. Auerbach and B. E. Larson, Phys. Rev. B **43**, 7800 (1991).
- E. Dagotto and J. Riera, Phys. Rev. Lett. **70**, 682 (1993).
- [6] W. O. Putikka, M. U. Luchini, and T. M. Rice, Phys. Rev. Lett. **68**, 538 (1992).
- [7] A. Moreo and D. J. Scalapino, Phys. Rev. B **43**, 8221 (1991); A. Moreo, D. J. Scalapino, and E. Dagotto, Phys. Rev. B **43**, 11442 (1991).
- [8] P. G. J. van Dongen, Phys. Rev. Lett. **74**, 182 (1995).
- [9] E. H. Lieb, Phys. Rev. Lett. **62**, 1201 (1989).
- [10] K. Kubo and T. Kishi, Phys. Rev. B **41**, 4866 (1990).
- [11] H. Shiba, Prog. Theor. Phys. **48**, 2171 (1972).
- [12] S. Q. Shen, and Z. M. Qiu, Phys. Rev. Lett. **71**, 4238 (1993).
- [13] C. N. Yang, Rev. Mod. Phys. **34**, 694 (1962); G. L. Sewell, J. Stat. Phys. **61**, 415 (1990); H. T. Nieh, G. Su, and B. H. Zhao, Phys. Rev. B **51**, 3760 (1995); C. Au and B. H. Zhao, Phys. Lett. A **209**, 235 (1995).